

Date: _____

Adult Patient: _____

PA: _____

Member #: _____

CM: _____

Admission Date _____

Behavioral Health Checklist for ASAM Criteria (3rd Edition)

Rehab	Level 1 OMT	Level 2.1	Level 2.5	Level 3.1	Levels 3.3, 3.5	Level 3.7	Level 4
Withdrawal Management	Level 2-WM	Level 2-WM				Level 3.7-WM	Level 4- WM
Description of Services	<p align="center">Outpatient (<9 hours/week) Or Amb Detox without extended on-site monitoring</p>	<p align="center">Intensive Outpatient (≥ 9 hours/week) Or Amb Detox with extended on-site monitoring</p>	<p align="center">Partial Hospitalization Programs</p>	<p align="center">Clinically Managed, Low Intensity Residential Services, e.g. Halfway House (professional staff on- site 24 hrs, or as req. by licensing regulations)</p>	<p align="center">(3.3)Clinically Managed, Population- Specific, High-Intensity Residential Services (Adult) (3.5) Clinically Managed High- Intensity Residential Services (Adult) AND (3.5) Clinically Managed Medium- Intensity Residential Services (Adolescent)</p>	<p align="center">Medically Monitored Intensive Inpatient (Adult) AND Medically Monitored High-Intensity Inpatient Services (Adolescent) Treatment (freestanding detox or special unit in psych hosp)</p>	<p align="center">Medically Managed Intensive Inpatient (Acute Inpatient Hosp)</p>
Admission specifications	<p align="center"><i>need to meet all Dimensions at this level</i></p>	<p align="center"><i>Advisable if meets Dimensions 2&3 at this level(if applicable), and one of Dimensions 4,5, & 6 at this level</i></p>	<p align="center"><i>Advisable if meets Dimension 2 at this level(if applicable), and one of Dimensions 4,5, & 6 at this level</i></p>	<p align="center"><i>need to meet all Dimensions at this level</i></p>	<p align="center"><i>need to meet all Dimensions at this level for III.3 and III.5; also programs need to be DDC or DDE based on Dimension 3</i></p>	<p align="center"><i>need to meet this level in 2 dimensions, at least one of which is Dimension 1,2, or 3.</i></p>	<p align="center"><i>need to meet at this level one of Dimensions 1, 2, or 3 at a minimum</i></p>
1 Acute Intoxication and/or Withdrawal Potential	<p>No withdrawal needs or needs can be safely managed at this level.</p> <p align="center"><input type="checkbox"/></p>	<p>No withdrawal needs or needs can be safely managed at this level.</p> <p align="center"><input type="checkbox"/></p>	<p>Moderate risk of severe withdrawal outside program, and can be safely managed with several hours/day of medical monitoring.</p> <p align="center"><input type="checkbox"/></p>	<p>No signs or symptoms of withdrawal, or withdrawal that can be managed at low intensity outpatient level</p> <p align="center"><input type="checkbox"/></p>	<p>Minimal risk of severe withdrawal, or if withdrawal is present, can be managed at a III.2- WM (social setting detox) level</p> <p align="center"><input type="checkbox"/></p>	<p>Experiencing severe withdrawal or severe withdrawal is imminent, but manageable in subacute detox or freestanding detox center, physician readily available within first 24 hours, and then 7 days/wk as needed</p> <p align="center"><input type="checkbox"/></p>	<p>Severe withdrawal risk complicated withdrawal requiring hourly monitoring (acute care inpatient, physician on-site daily and available 24-hr/day)</p> <p align="center"><input type="checkbox"/></p>
2 Biomedical Conditions and Complications	<p>None or sufficiently stable</p> <p align="center"><input type="checkbox"/></p>	<p>If present, are stable or are receiving concurrent attention and won't interfere with treatment.</p> <p align="center"><input type="checkbox"/></p>	<p>Biomedical conditions won't interfere with treatment, but would distract from recovery without this Level of management.</p> <p align="center"><input type="checkbox"/></p>	<p>Stable and no medical monitoring needed, <i>or</i> doesn't warrant inpatient, but needs medical monitoring.</p> <p align="center"><input type="checkbox"/></p>	<p>If present, stable and can self-administer meds <i>or</i> severe enough to warrant medical monitoring but not in need of inpatient treatment</p> <p align="center"><input type="checkbox"/></p>	<p>Serious risk to physical health or requires 24-hour nursing and medical monitoring, but not full resources of an acute care hospital</p> <p align="center"><input type="checkbox"/></p>	<p>Requires inpatient diagnosis and treatment (24-hour medical, nursing care)</p> <p align="center"><input type="checkbox"/></p>

Date: _____

Adult Patient: _____

PA: _____

Member #: _____

CM: _____

Admission Date _____

Rehab	Level 1 OMT	Level 2.1	Level 2.5	Level 3.1	Levels 3.3, 3.5	Level 3.7	Level 4
Withdrawal Management	Level 2-WM	Level 2-WM				Level 3.7-WM	Level 4- WM
3 Emotional/ Behavioral or Cognitive Conditions and Complications	None or very stable (cognitively able to participate and no risk of harm) <input type="checkbox"/>	If present, mild severity and needs outpatient monitoring to minimize distractions from recovery. <input type="checkbox"/>	Problems here not required for this Level; if present, program needs to be DDC or DDE, and problems do not require a 24- hour level of care. <input type="checkbox"/>	Stable or if distracting, can respond to the level of 24-hour structure in this program. <input type="checkbox"/>	Moderate severity needing a 24-hour structured setting for cognitive deficits; repeated inability to control impulses, personality disorder requires high structure to shape behavior <input type="checkbox"/>	Unstable BH condition that requires 24-hour medical monitoring and/or has violent behavior posing imminent danger <input type="checkbox"/>	Dual diagnosis, severe problems requiring 24-hour psychiatric care with concomitant addiction treatment <input type="checkbox"/>
4 Readiness to Change	Willing to cooperate or is ambivalent and needs motivating and monitoring strategies <input type="checkbox"/>	Resistance high enough to require structured program but not so high as to render outpatient treatment ineffective. <input type="checkbox"/>	This degree of structure required due to failure of motivational interventions at lesser levels, or poor impulse control and follow through with less intensive programs. <input type="checkbox"/>	Ready to change and cooperate at this level, <i>or</i> externalizes problems and needs this level of structure, motivating and support. <input type="checkbox"/>	Limited readiness to change; despite serious consequences, doesn't link addiction and life problems; without this structure poses a danger to self or others with continued use.; perspective impairs ability to change without 24 hour structure. <input type="checkbox"/>	Resistance high and impulse control poor despite negative consequences; needs intensive motivation strategies available only in 24-hour structure; marked opposition to treatment with dangerous consequences if not engaged in treatment <input type="checkbox"/>	Problems in this dimension do not qualify for Level IV treatment <input type="checkbox"/>
5 Relapse, Continued Use or Continued Problem Potential	Able to maintain abstinence and recovery goals or achieve awareness of a SA problem with minimal support <input type="checkbox"/>	Intensification of symptoms despite active participation (modification of plan) in Level I, and high likelihood of relapse without close monitoring and support <input type="checkbox"/>	Despite active involvement and modification of treatment plans at lower levels, continues to be experiencing deterioration in functioning/symptoms. <input type="checkbox"/>	Is at high risk for imminent relapse with dangerous consequences unless provided this 24-hour structure and support <i>or</i> needs this support to transition into community. <input type="checkbox"/>	Doesn't recognize triggers, unable to control use, in danger of relapse without close 24-hour monitoring and structured treatment <input type="checkbox"/>	Unable to control use despite active participation in less intensive care and needs 24-hour structure <input type="checkbox"/>	Problems in this dimension do not qualify for Level IV treatment <input type="checkbox"/>
6 Recovery / Living Environment	Supportive recovery environment or willingness to obtain such or supports need professional interventions. <input type="checkbox"/>	Lacks social contacts or social contacts aren't conducive to recovery, but with structure or support, the patient can cope <input type="checkbox"/>	Environment is unsupportive but with structure or support and relief from the home environment, the patient can cope <input type="checkbox"/>	Has a using, unsupportive, dangerous, or victimizing social network, <i>or</i> lacks a social network, requiring this level of 24-hour support. <input type="checkbox"/>	Moderately high risk environment, lacks appropriate social contacts, is in danger of victimization <input type="checkbox"/>	Environment is dangerous for recovery, patient lacks skills to cope outside highly structured 24-hour setting <input type="checkbox"/>	Problems in this dimension do not qualify for Level IV treatment <input type="checkbox"/>

Date: _____

Adult Patient: _____

PA: _____

Member #: _____

CM: _____

Admission Date _____

Rehab	Level 1 OMT	Level 2.1	Level 2.5	Level 3.1	Levels 3.3, 3.5	Level 3.7	Level 4
Withdrawal Management	Level 2-WM	Level 2-WM				Level 3.7-WM	Level 4- WM

The ratings of the above dimensions indicate that ASAM criteria are met for _____ Level of care.